

The Role of Neuroplasticity-Based Interventions in Improving Cognitive Functioning

Dr. Vipul Vithal¹

Dr. Mansi Kapoor²

¹Assistant Professor, Faculty of Education, Desh Bhagat University, Mandi Gobindgarh

²Assistant Professor, Faculty of Education, Desh Bhagat University, Mandi Gobindgarh

Abstract

Neuroplasticity is the capacity of the brain and nervous system to modify their structure, organization, and functional connectivity in response to learning, experience, environmental stimulation, and injury. The recognition that the brain retains considerable plasticity throughout adulthood has significantly influenced contemporary approaches to cognitive enhancement and rehabilitation. Neuroplasticity-based interventions—including cognitive training, physical exercise, mindfulness-based practices, computerized cognitive interventions, skill learning, and cognitive rehabilitation—seek to stimulate adaptive changes in neural networks and consequently improve cognitive functioning. This review examines the theoretical foundations, neurobiological mechanisms, empirical evidence, applications, limitations, and future directions associated with these interventions. A review of systematic reviews, meta-analyses, and clinical research indicates that neuroplasticity-based interventions can produce improvements in selected cognitive domains, although effects vary according to intervention type, population, intensity, duration, and outcome measure. Working-memory training demonstrates substantial improvements on trained tasks and moderate near-transfer effects, while evidence for far transfer to unrelated cognitive abilities is considerably weaker. Physical exercise appears to support neuroplasticity through mechanisms involving brain-derived neurotrophic factor (BDNF), cerebral circulation, vascular health, synaptic functioning, and hippocampal plasticity. Recent meta-analytic evidence indicates that exercise can increase resting BDNF levels in older adults. Multicomponent exercise has also been associated with improvements in cognition, attention, and executive functioning among older adults with mild cognitive impairment or dementia. Mindfulness-based interventions show small but meaningful improvements in several aspects of cognitive functioning, although their effects are not uniform across cognitive domains. Cognitive rehabilitation demonstrates particular value following acquired brain injury, stroke, and traumatic brain injury, especially when interventions are individualized and functionally oriented. Overall, neuroplasticity-based interventions represent promising non-pharmacological approaches for improving and maintaining cognitive functioning. However, the evidence does not support the assumption that all interventions produce generalized increases in intelligence or cognition. Future research

should emphasize personalized intervention design, multimodal approaches, longer follow-up periods, standardized cognitive outcomes, and integration of behavioral, neuroimaging, and biological measures.

Keywords: *neuroplasticity, cognitive functioning, cognitive training, working memory, physical exercise, mindfulness, cognitive rehabilitation, BDNF, brain health*

1. Introduction

Cognitive functioning refers to the collection of mental processes that enable individuals to acquire, process, store, retrieve, and use information. Major cognitive domains include attention, memory, executive functioning, processing speed, reasoning, language, problem-solving, and cognitive flexibility. These abilities are fundamental to academic achievement, occupational performance, social interaction, independent living, and psychological well-being.

Cognitive functioning is not static. It changes throughout the lifespan as a consequence of biological development, learning, environmental experience, lifestyle, aging, illness, and neurological injury. Although some aspects of cognition may decline with age or neurological disease, contemporary neuroscience has demonstrated that the adult brain retains the capacity for structural and functional modification. This capacity is generally described as neuroplasticity.

Neuroplasticity involves changes at multiple levels of the nervous system. These include alterations in synaptic strength, dendritic structure, neuronal connectivity, cortical representation, and functional organization of brain networks. Learning and repeated experience can therefore modify the efficiency with which neural systems process information. The concept of neuroplasticity has important implications for cognitive psychology, education, rehabilitation, gerontology, and neuroscience. If the brain can reorganize itself in response to experience, appropriately designed interventions may be capable of strengthening cognitive abilities and supporting recovery following neurological damage. Consequently, researchers have developed interventions designed to stimulate neuroplastic change. These include working-memory training, computerized cognitive training, aerobic and resistance exercise, mindfulness meditation, skill learning, environmental enrichment, and cognitive rehabilitation. These approaches differ considerably in their theoretical foundations and methods, but they share the assumption that repeated and meaningful engagement can modify neural functioning. The purpose of this review is to examine the role of neuroplasticity-based interventions in improving cognitive functioning. Particular attention is given to the mechanisms underlying neuroplasticity, major intervention approaches, evidence for cognitive improvement, applications across different populations, limitations of existing research, and directions for future investigation.

2. Conceptualizing Neuroplasticity

Neuroplasticity can be defined as the nervous system's capacity to alter its structure and function in response to internal or external demands. The concept encompasses several processes, including synaptic plasticity, structural remodeling, cortical reorganization, neurogenesis, and changes in functional connectivity. Historically, the adult human brain was often considered relatively fixed after development. This view has been replaced by evidence demonstrating that learning, environmental exposure, physical activity, and rehabilitation can alter neural organization throughout adulthood. Neuroplasticity is not inherently positive. Neural systems can adapt in response to both beneficial and harmful experiences. Repeated practice of a useful skill may strengthen adaptive neural pathways, whereas chronic stress, inactivity, substance exposure, or maladaptive behaviors may contribute to less desirable patterns of neural functioning.

Therefore, the central question for intervention research is not simply whether the brain is plastic, but how plasticity can be directed toward adaptive cognitive outcomes.

3. Neurobiological Foundations of Neuroplasticity

3.1 Synaptic Plasticity

Synaptic plasticity refers to changes in the strength and efficiency of communication between neurons. Long-term potentiation (LTP) and long-term depression (LTD) are two important mechanisms.

Repeated activation of neural pathways can strengthen synaptic connections, while reduced activity can weaken them. These processes provide a biological foundation for learning and memory.

3.2 Dendritic Remodeling

Learning and environmental stimulation can influence dendritic branching and spine density. Dendrites receive signals from other neurons, and changes in dendritic structure can affect the efficiency of neural communication.

3.3 Functional Connectivity

Cognitive processes depend on communication between distributed brain networks. Neuroplasticity can modify the functional relationships between brain regions, potentially making cognitive processing more efficient. For example, cognitive training can produce changes in frontoparietal systems involved in executive functioning and working memory. A systematic review and meta-analysis of working-memory updating training found consistent neural changes alongside behavioral improvements.

3.4 Brain-Derived Neurotrophic Factor

BDNF is a neurotrophin involved in neuronal survival, synaptic plasticity, learning, and memory. Physical exercise is particularly relevant to this mechanism. A 2025 systematic review and meta-analysis of 35 randomized controlled trials found that exercise training significantly increased resting BDNF levels in older adults, with effects observed across aerobic, resistance, and combined exercise modalities. These findings provide biological support for the hypothesis that exercise can create an environment conducive to neuroplasticity.

4. Neuroplasticity Across the Lifespan

Neuroplasticity occurs throughout life, although its characteristics may differ across developmental stages. During childhood and adolescence, the brain undergoes extensive structural development and network specialization. Learning experiences during these periods can produce substantial neural adaptations. During adulthood, neuroplasticity continues through learning, occupational experience, physical activity, and environmental stimulation. Although the rate of some forms of plasticity may decrease with age, older adults continue to demonstrate neural adaptation. Age-related cognitive decline is therefore not necessarily synonymous with irreversible cognitive deterioration. Lifestyle and behavioral factors can influence cognitive trajectories. The existence of neuroplasticity in later life provides an important theoretical foundation for interventions aimed at maintaining cognitive health among older adults.

5. Major Neuroplasticity-Based Interventions

5.1 Cognitive Training

Cognitive training consists of repeated exercises designed to strengthen specific cognitive abilities. Common targets include:

- Working memory
- Attention
- Processing speed
- Reasoning
- Inhibitory control
- Cognitive flexibility
- Problem-solving

Working-memory updating has received considerable research attention.

Pappa et al. (2020) conducted a systematic review and meta-analysis examining working-memory updating training. The authors reported a large training effect and a moderate near-transfer effect, but the effect on far-transfer tasks involving unrelated cognitive domains was not statistically significant.

This distinction between training gains, near transfer, and far transfer is central to evaluating cognitive training.

Training Gain

A training gain occurs when performance improves on the task directly practiced.

Near Transfer

Near transfer occurs when improvements generalize to tasks that are similar to the trained task.

Far Transfer

Far transfer refers to improvements in relatively unrelated abilities, such as general intelligence or academic performance.

Current evidence suggests that training gains are generally stronger than far-transfer effects. Therefore, cognitive training should not automatically be interpreted as a method for producing generalized increases in intellectual ability.

6. Physical Exercise and Neuroplasticity

Physical exercise is one of the most extensively studied behavioral approaches to brain health.

Exercise may promote neuroplasticity through several pathways:

- Increased cerebral blood flow.
- Increased BDNF.
- Improved cardiovascular health.
- Enhanced synaptic functioning.
- Reduced inflammation.
- Improved metabolic health.
- Support for hippocampal functioning.
- Improved mood and stress regulation.

A systematic review and meta-analysis involving older adults with mild cognitive impairment or dementia found that multicomponent exercise significantly improved cognition, attention, and executive functioning, although significant improvement in memory was not demonstrated.

Recent evidence also indicates that exercise training increases BDNF levels in older adults. Gholami et al. (2025) reported significant increases following aerobic, resistance, and combined exercise programs. These findings suggest that physical exercise may influence cognition through both direct neural mechanisms and indirect effects on vascular, metabolic, and psychological health.

7. Mindfulness-Based Interventions

Mindfulness involves maintaining attention on present-moment experiences while adopting an attitude of openness and nonjudgment. Mindfulness-based interventions have been investigated for:

- Attention
- Working memory
- Executive functioning
- Cognitive flexibility
- Inhibitory control
- Emotional regulation

Zainal and Newman (2023) conducted a meta-analysis of 111 randomized controlled trials and found that mindfulness interventions produced improvements across several cognitive domains, although effects varied by outcome and were generally modest. The potential neuroplastic mechanisms of mindfulness include changes in attentional networks, emotion-regulation systems, and functional connectivity. However, mindfulness should not be presented as a universal cognitive enhancement method. Evidence indicates domain-specific effects, and improvements may be influenced by intervention duration, participant characteristics, adherence, and the quality of the control condition.

8. Cognitive Rehabilitation

Cognitive rehabilitation is particularly important for individuals whose cognitive abilities have been affected by neurological injury or disease.

It may be used following:

- Stroke
- Traumatic brain injury
- Brain tumor
- Neurological disease
- Other acquired brain injuries

Cognitive rehabilitation may use restorative techniques, compensatory strategies, metacognitive approaches, and functional training. Cicerone et al. (2019) reviewed a large body of evidence concerning cognitive rehabilitation following traumatic brain injury and stroke. Their review supported evidence-based recommendations for interventions addressing attention, memory, executive functioning, language, social communication, and broader functional disability. Similarly, a meta-analysis of cognitive training following chronic traumatic brain injury found a small but statistically significant effect on overall cognition and a moderate effect on overall functional outcomes. These findings demonstrate that

neuroplasticity-based approaches have applications beyond healthy cognitive enhancement and may contribute to functional recovery following neurological injury.

9. Neuroplasticity and Specific Cognitive Functions

9.1 Attention

Attention enables individuals to selectively process relevant information while filtering distractions. Cognitive training and rehabilitation can target sustained, selective, divided, and alternating attention. Evidence from acquired brain injury research indicates that cognitive rehabilitation can improve some forms of attention, although effects are not consistent across all attention domains or populations.

9.2 Working Memory

Working memory enables individuals to temporarily maintain and manipulate information.

Working-memory training is one of the most extensively researched neuroplasticity-based interventions. Evidence supports substantial improvements on trained tasks and moderate near-transfer effects, but generalized far transfer remains uncertain.

9.3 Long-Term Memory

Memory depends on complex interactions among the hippocampus, medial temporal structures, cortical networks, and executive systems. Exercise may support memory indirectly through vascular health, neurotrophic factors, and hippocampal functioning. However, evidence is not uniformly positive, particularly among populations with cognitive impairment.

9.4 Executive Function

Executive functioning includes:

- Planning
- Inhibition
- Cognitive flexibility
- Decision-making
- Problem-solving
- Monitoring

Exercise appears particularly promising for executive functioning. In older adults with mild cognitive impairment or dementia, multicomponent exercise has demonstrated improvements in executive functioning.

9.5 Processing Speed

Processing speed refers to how quickly an individual can perceive, interpret, and respond to information.

Cognitive and physical interventions may improve aspects of processing efficiency, although findings vary according to intervention and assessment method.

10. Comparative Overview of Major Interventions

Intervention	Primary Target	Proposed Mechanisms	Evidence
Cognitive training	Working memory, attention, reasoning	Synaptic strengthening, network efficiency	Strong training effects; limited far transfer
Aerobic exercise	Global cognition, executive function	BDNF, circulation, neurotrophic mechanisms	Generally positive
Resistance exercise	Executive and global cognition	BDNF, metabolic and vascular effects	Increasing evidence
Mindfulness	Attention, executive function, working memory	Network regulation, attentional control	Small-to-moderate effects
Cognitive rehabilitation	Functional cognition	Reorganization, compensatory networks	Strong clinical evidence for selected deficits
Skill learning	Task-specific cognition	Repeated neural activation	Strong task-specific plasticity
Multicomponent exercise	Global cognition	Combined physical and neurobiological mechanisms	Promising

11. Evidence from Systematic Reviews and Meta-Analyses

Systematic reviews and meta-analyses provide important evidence regarding the effectiveness of neuroplasticity-based interventions.

Cognitive Training

Pappa et al. (2020) found strong improvements in trained working-memory tasks and moderate near-transfer effects, but no statistically significant far-transfer effect.

Exercise

Wang et al. (2020) found significant improvements in cognition, attention, and executive functioning among community-dwelling older adults with mild cognitive impairment or dementia following multicomponent exercise.

Mindfulness

Zainal and Newman (2023) synthesized 111 randomized controlled trials and found that mindfulness-based interventions improved several cognitive outcomes, although effects were generally modest and domain-dependent.

Cognitive Rehabilitation

Cicerone et al. (2019) identified substantial evidence supporting cognitive rehabilitation approaches for attention, memory, executive functioning, communication, and functional disability following acquired brain injury.

Taken together, these findings indicate that neuroplasticity-based interventions are most defensible when their intended outcomes are clearly defined and closely matched to the intervention.

12. Applications in Education

The concept of neuroplasticity has important implications for education.

Learning itself represents a form of experience-dependent plasticity. Educational activities repeatedly activate cognitive and neural systems, potentially strengthening the networks underlying academic skills.

Neuroplasticity-informed educational practices may include:

- Retrieval practice
- Repeated meaningful practice
- Multisensory learning
- Physical activity
- Attention training
- Metacognitive strategies
- Mindfulness exercises
- Problem-based learning
- Skill-based learning

However, educators should avoid simplistic claims that any particular "brain training" program will dramatically increase intelligence.

A neuroplasticity-informed educational approach should instead emphasize the importance of sustained practice, feedback, meaningful learning, motivation, physical activity, adequate sleep, and supportive environments.

13. Applications in Older Adults

Population aging has increased interest in interventions capable of maintaining cognitive functioning.

Older adults may experience gradual changes in processing speed, memory, attention, and executive functioning. Neuroplasticity-based interventions may help maintain cognitive reserve and support healthy aging.

Exercise is particularly promising because it simultaneously affects cardiovascular health, metabolic functioning, mood, and neural mechanisms.

Recent evidence indicates that exercise can significantly increase BDNF in older adults, providing a plausible biological pathway linking physical activity with neuroplasticity.

Nevertheless, exercise should be understood as a component of a broader healthy-aging strategy rather than as a guaranteed method of preventing dementia.

14. Applications in Neurological Rehabilitation

Neuroplasticity has transformed neurological rehabilitation.

Following brain injury, the nervous system may reorganize through:

- Recruitment of alternative neural networks.
- Strengthening of surviving pathways.
- Reorganization of cortical representations.
- Development of compensatory strategies.
- Repeated task-specific practice.

The principle of use-dependent plasticity is particularly important. Functional improvement requires meaningful practice of the behaviors that individuals need to regain.

Cognitive rehabilitation therefore emphasizes repetition, task specificity, feedback, goal setting, and functional relevance.

Evidence-based reviews support cognitive rehabilitation following stroke and traumatic brain injury, particularly for attention, memory, executive functioning, language, and functional independence.

15. Role of Intensity, Duration, and Repetition

Neuroplastic change does not occur simply because an intervention is provided. The characteristics of the intervention are important.

Important variables include:

- Frequency
- Duration
- Intensity
- Repetition
- Task difficulty
- Feedback
- Motivation
- Individual baseline ability

- Environmental context

Exercise research illustrates this point. Gholami et al. (2025) found that exercise interventions lasting approximately 12 weeks and involving moderate-to-vigorous or vigorous intensity and three to four sessions per week produced relatively larger BDNF effects, although intervention-specific conclusions should be interpreted cautiously.

This supports the broader principle that neuroplastic interventions should be sufficiently intensive and sustained to provide repeated stimulation.

16. Individual Differences and Personalized Neuroplasticity

Not all individuals respond to interventions in the same way.

Potential moderators include:

- Age
- Baseline cognitive ability
- Neurological status
- Motivation
- Education
- Genetics
- Physical fitness
- Sleep
- Stress
- Intervention adherence
- Social environment

This variation has led to increasing interest in personalized neurorehabilitation and cognitive training.

Future interventions may use baseline cognitive assessments, neuroimaging, physiological measures, and behavioral profiles to determine which intervention is most appropriate for a particular individual.

Personalization is particularly important in rehabilitation because two individuals with the same diagnostic label may have very different patterns of cognitive impairment.

17. Critical Evaluation of the Evidence

Although the evidence supporting neuroplasticity-based interventions is promising, several limitations must be recognized.

17.1 Transfer Effects

One of the most important limitations is the distinction between improvement on a trained task and improvement in everyday cognitive functioning.

Cognitive training may make individuals better at the specific activities they practice without producing substantial changes in broader cognitive abilities.

The working-memory literature provides a clear example: strong training effects and moderate near-transfer effects coexist with limited evidence for far transfer.

17.2 Methodological Heterogeneity

Studies vary substantially in:

- Sample size
- Intervention duration
- Intervention intensity
- Control conditions
- Cognitive assessments
- Participant characteristics

This heterogeneity makes it difficult to establish a single optimal intervention protocol.

17.3 Short Follow-Up Periods

Many studies measure cognition immediately after intervention. Less is known about whether improvements remain months or years later.

Long-term follow-up is therefore essential.

17.4 Publication Bias

Positive findings may be more likely to be published than null findings. This can lead to an overestimation of intervention effectiveness.

17.5 Brain Changes Versus Behavioral Changes

Neuroimaging evidence of altered brain activity does not necessarily mean that meaningful improvements in everyday cognitive functioning have occurred.

A complete evaluation should therefore integrate:

- Neural measures.
- Standardized cognitive tests.
- Functional outcomes.
- Real-world behavior.

- Long-term follow-up.

18. Ethical Considerations

Neuroplasticity-based interventions should be implemented ethically.

Researchers and practitioners should avoid exaggerated claims regarding cognitive enhancement. Terms such as "rewiring the brain" can be misleading when used without scientific qualification.

Participants should receive accurate information about:

- Expected benefits
- Possible limitations
- Duration of intervention
- Evidence quality
- Alternative approaches

In clinical rehabilitation, interventions should be based on individual goals and evidence-based practice rather than commercial claims.

19. Future Directions

Future research should move toward more sophisticated models of neuroplasticity-based intervention.

19.1 Multimodal Interventions

Combining exercise, cognitive training, mindfulness, and social engagement may provide complementary forms of stimulation.

19.2 Precision Intervention

Researchers should identify individual characteristics that predict treatment response.

19.3 Digital Technologies

Artificial intelligence, virtual reality, adaptive cognitive training, and wearable technologies may enable interventions to adjust automatically according to performance.

19.4 Neuroimaging-Guided Rehabilitation

Functional MRI, diffusion imaging, EEG, and other measures may help identify neural changes associated with successful rehabilitation.

19.5 Longitudinal Research

Future studies should follow participants for months or years rather than relying primarily on immediate post-intervention outcomes.

19.6 Real-World Outcomes

Researchers should examine whether cognitive improvements translate into:

- Academic achievement
- Employment
- Independent living
- Social functioning
- Quality of life
- Reduced caregiver burden

Such outcomes are ultimately more meaningful than performance on laboratory tasks alone.

20. Implications for Researchers and Practitioners

The current evidence suggests several practical principles.

First, interventions should be specific and goal-oriented. A person with working-memory difficulties may benefit from working-memory exercises, whereas an individual recovering from stroke may require a broader rehabilitation program.

Second, interventions should be repeated and sufficiently intensive.

Third, cognitive interventions should ideally be integrated with healthy lifestyle practices, particularly physical activity, adequate sleep, and social engagement.

Fourth, progress should be measured using standardized assessments and functional outcomes.

Finally, practitioners should communicate realistic expectations. Neuroplasticity provides an important biological foundation for change, but it does not imply unlimited cognitive enhancement.

21. Conclusion

Neuroplasticity represents one of the most important concepts in contemporary neuroscience and cognitive rehabilitation. The recognition that the brain can reorganize itself throughout life provides a strong theoretical foundation for interventions designed to improve cognitive functioning.

The evidence reviewed in this paper indicates that cognitive training, physical exercise, mindfulness-based interventions, and cognitive rehabilitation can produce meaningful improvements in selected cognitive domains. Cognitive training is particularly effective for trained abilities and related skills, although far-transfer effects remain limited. Physical exercise provides broad support for brain health and may promote neuroplasticity through BDNF, vascular, metabolic, and synaptic mechanisms. Mindfulness appears to produce modest improvements in several cognitive domains, particularly executive and attentional processes.

Cognitive rehabilitation has demonstrated important clinical benefits following stroke and traumatic brain injury.

However, neuroplasticity should not be interpreted as evidence that cognitive abilities can be improved without limits. The effects of interventions depend on individual characteristics, intervention intensity, duration, adherence, task specificity, and outcome measurement. In particular, improvements in laboratory-based cognitive tasks do not necessarily translate into broad improvements in everyday functioning.

The future of neuroplasticity-based intervention is therefore likely to involve personalized, multimodal, evidence-based approaches that combine cognitive stimulation, physical activity, mindfulness, rehabilitation, and supportive environments. Integration of behavioral assessments with neuroimaging and biological measures may further clarify how neural changes translate into meaningful cognitive and functional improvements.

Ultimately, neuroplasticity offers a scientifically grounded explanation for the possibility of cognitive change across the lifespan. Its greatest value lies not in promising unlimited brain enhancement, but in providing a framework for designing interventions that help individuals learn, adapt, recover, and maintain cognitive functioning.

References

- Cicerone, K. D., Goldin, Y., Ganci, K., Rosenbaum, A., Wethe, J. V., Langenbahn, D. M., Malec, J. F., Bergquist, T. F., Kingsley, K., Nagele, D., Trexler, L., Fraas, M., Bogdanova, Y., & Harley, J. P. (2019). Evidence-based cognitive rehabilitation: Systematic review of the literature from 2009 through 2014. *Archives of Physical Medicine and Rehabilitation*, 100(8), 1515–1533. <https://doi.org/10.1016/j.apmr.2019.02.011>
- Gholami, F., Mesrabadi, J., Iranpour, M., & Donyaei, A. (2025). Exercise training alters resting brain-derived neurotrophic factor concentration in older adults: A systematic review with meta-analysis of randomized-controlled trials. *Experimental Gerontology*, 199, 112658. <https://doi.org/10.1016/j.exger.2024.112658>
- Pappa, K., Biswas, V., Flegal, K. E., Evans, J. J., & Baylan, S. (2020). Working memory updating training promotes plasticity and behavioural gains: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, 118, 209–235. <https://doi.org/10.1016/j.neubiorev.2020.07.027>
- Wang, X., Wang, H., Ye, Z., Ding, G., Li, F., Ma, J., & Hua, W. (2020). The neurocognitive and BDNF changes of multicomponent exercise for community-dwelling older adults with mild cognitive impairment or dementia: A systematic review and meta-analysis. *Aging*, 12(6), 4907–4917. <https://doi.org/10.18632/aging.102918>
- Zainal, N. H., & Newman, M. G. (2023). Mindfulness enhances cognitive functioning: A meta-analysis of 111 randomized controlled trials. *Health Psychology Review*, 18(2), 369–395. <https://doi.org/10.1080/17437199.2023.2248222>



- Mewborn, C. M., Lindbergh, C. A., & Stephen Miller, L. (2017). Cognitive interventions for cognitive aging: The role of intellectual engagement and social engagement. *Psychology and Aging*, 32(1), 85–98.
- Reis, J., Schambra, H. M., Cohen, L. G., Buch, E. R., Fritsch, B., Zarahn, E., & Krakauer, J. W. (2009). Noninvasive cortical stimulation enhances motor skill acquisition over multiple days through an effect on consolidation. *Proceedings of the National Academy of Sciences*, 106(5), 1590–1595. <https://doi.org/10.1073/pnas.0805413106>
- Voss, M. W., Prakash, R. S., Erickson, K. I., Boot, W. R., Burdette, J. H., Kramer, A. F., & Colcombe, S. J. (2010). Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Frontiers in Aging Neuroscience*, 2, 32. <https://doi.org/10.3389/fnagi.2010.00032>
- Cicerone, K. D., Langenbahn, D. M., Braden, C., Malec, J. F., Kalmar, K., Fraas, M., Felicetti, T., Laatsch, L., Harley, J. P., Bergquist, T., Azulay, J., Cantor, J., & Ashman, T. (2011). Evidence-based cognitive rehabilitation: Updated review of the literature from 2003 through 2008. *Archives of Physical Medicine and Rehabilitation*, 92(4), 519–530. <https://doi.org/10.1016/j.apmr.2010.11.015>
- Zelinski, E. M., & Reyes, R. (2009). Cognitive benefits of computer games for older adults: A review. *Gerontechnology*, 8(4), 218–229.